25

## CLAIMS

97 a17:

- 1. Adevice for iontophoresis supplying a drug to transdermal or transmcosal, comprising: first means for detecting a capacitance stored in the transdermal or the transmcosal; and second means for determining a conduction state of current into the transdermal or the transmcosal based on the output detected by the first means.
- The device for ion tophoresis according to Claim 1, wherein
   the first means is a detection circuit for a reactive current flowing through the transdermal or the transmcosal.
  - 3. The device for iontophoresis according to Claim 1, wherein the first means is a detection circuit for a residual voltage developed in the transdermal or the transmcosal.
- 4. A method for determining an operation of an iontophoresis apparatus, wherein a capacitance stored in transdermal or transmcosal is detected to determine a conduction state of current flowing into the transdermal or the transmcosal.
- 5. The method for detecting an operation of an iontophoresis
  apparatus according to Claim 4, wherein the detection of the
  capacitance is carried out by detecting a reactive current
  flowing through the transdermal or the transmcosal.
  - 6. The method for detecting an operation of an iontophoresis apparatus according to Claim 4, wherein the detection of the capacitance is carried out by detecting a residual voltage developed in the transdermal or the transmosal.
  - 7. An iontophoresis apparatus comprising: a preparation for

qub C1

5

iontophoressis, holding a drug; and a device for iontophoresis having means for generating an electrical output to supply a drug from the preparation into transdermal or transmcosal and means for detecting a capacitance stored in the transdermal or the transmcosal to determine a conduction state of a current flowing into the transdermal or the transmcosal.